cient ra iative transfer mo elin of scatterin polariation

P. Benedusi[1], S. Riva[1], G. Janett[1], and L. Beluzzi[1]

[1] *s i u o ricerche solari l o e ele acc acul o n orma ics niersi ella iera i aliana ocarno i erlan*

[2] *uler ns i u e acul o n orma ics niersi ella iera i aliana u ano i erlan*

contact e-mail: pietro.benedusi@usi.ch

e present an e cient and massively parallel solution strategy for the transfer problem of polari ed radiation, for a 3D stationary medium out of local thermodynamic equilibrium. Scattering processes are included accounting for partial frequency redistribution effects. Such a setting is one of the most challenging ones in radiative transfer modeling. The problem is formulated for a two-level atomic model, which allows lineari ation. The discrete ordinate method alongside an exponential integrator are used for discreti ation. E cient solution is obtained with a Krylov method equipped with a tailored physics-based preconditioner. A matrix-free approach results in a lightweight implementation, suited for tackling large problems. ear-optimal strong and weak scalability are obtained with two complementary decompositions of the computational domain. The presented approach made it possible to perform simulations for the Ca I line at 422 Å with more than 40 billions degrees of freedom in few hours on massively parallel machines, always converging in a few iterations for the proposed tests.